Tropical infectious disease research

Steve Meshnick

Malawi

- Malaria during pregnancy
 - Intermittent Preventive Therapy (IPT)
 - Treatments for HIV-malaria coinfections
- HIV mother-to-child transmission
 - Syphilis coinfections increase risk ~3x
 - Maternal-fetal microtransfusions
 - CHAVI 009
- Archived chloroquine resistance
- > U of Malawi School of Public Health

Is chloroquine resistance really gone from Malawi?

The NEW ENGLAND JOURNAL of MEDICINE

SETS S1.100 SD 101 381.2

MOVEMBER 9, 2006

704, 355 HO. 19


Return of Chloroquine Antimalarial Efficacy in Malawi


Miriam K. Laufer, M.D., Phillip C. Thesing, D.O., Nicole D. Eddington, M.S., Rhoda Masonga, Fraction K. Dzinjalamala, Ph.D., Shannon L. Talala, Ph.D., Teme E. Taylor, D.O., and Christopher V. Plowe, M.D., M.P.H.

CVIET Heteroduplex
Wild Type
Heteroduplex
Positive
Samples

Cambodia/Thailand

- Monitor drug resistance
- Change drug policy





DR Congo

- Malaria during pregnancy
 - Prompt treatment minimalizes adverse effects
 - Malnutrition is an effect measure modifier
 - Development of new IPT agents
- > Human African Trypanosomiasis
 - Epidemiology
 - Drug access

Historic HAT Foci

